

Consensus Findings on Meibomian Gland Dysfunction Published

ScienceDaily (Mar. 30, 2011) — The first global consensus report on meibomian gland dysfunction -- a major cause of lid disease and evaporative dry eye -- has been published in a special issue of the *Investigative Ophthalmology & Visual Science* (IOVS) journal. The report is the result of findings from a two-year-long workshop composed of more than 50 leading clinical and basic research experts from around the world.

The workshop participants used an evidence-based approach to develop a worldwide definition: meibomian gland dysfunction (MGD) is a chronic, diffuse abnormality of the meibomian glands, commonly characterized by terminal duct obstruction and/or qualitative/quantitative changes in the glandular secretion. This may result in alteration of the tear film, symptoms of eye irritation, clinically apparent inflammation, and ocular surface disease.

Using the same methodology, the participants developed a universal classification system -- based on pathophysiology, rather than anatomical changes or the severity of disease -- to meet the needs of clinicians and researchers alike. The consensus paper further proposes recommendations for diagnosing MGD and MGD-related disorders and presents a sequence of diagnostic tests to be performed in an order that will minimize the extent to which one test influences those that follow.

Also included in the report are recommendations for the evaluation and grading of the severity of MGD, management of and therapy for the disease and norms for clinical trials designed to evaluate pharmaceutical interventions for treatment.

The International Workshop on Meibomian Gland Dysfunction was conducted by the Tear Film & Ocular Surface Society (TFOS). While the breadth and depth of the consensus findings are expected to have a far-reaching impact on the clinical care of patients, the group of experts concur that additional research be conducted to further study other aspects of MGD. These include its association with dry eye disease and standardized and validated ways to identify symptoms and signs of MGD.

Email or share this

| [More](#)

story:

Story Source:

The above story is reprinted (with editorial adaptations by *ScienceDaily* staff) from materials provided by [Association for Research in Vision and Ophthalmology](#), via [EurekAlert!](#), a service of AAAS.

Journal Reference:

1. K. K. Nichols, G. N. Foulks, A. J. Bron, B. J. Glasgow, M. Dogru, K. Tsubota, M. A. Lemp, D. A. Sullivan. **The International Workshop on Meibomian Gland Dysfunction: Executive Summary.** *Investigative Ophthalmology & Visual Science*, 2011; 52 (4): 1922 DOI: [10.1167/iovs.10-6997a](https://doi.org/10.1167/iovs.10-6997a)

Need to cite this story in your essay, paper, or report? Use one of the following formats:

APA

MLA

Association for Research in Vision and Ophthalmology (2011, March 30). Consensus findings on meibomian gland dysfunction published. *ScienceDaily*. Retrieved June 7, 2011, from <http://www.sciencedaily.com/releases/2011/03/110331094611.htm>

Note: If no author is given, the source is cited instead.

Disclaimer: *This article is not intended to provide medical advice, diagnosis or treatment. Views expressed here do not necessarily reflect those of ScienceDaily or its staff.*